



else go to 2

2. compute  $x^*$  - median of all  $x$ -coordinates of points in  $S$

$S_1 - p \in S : x(p) < x^*$

$S_2 - p \in S : x(p) > x^*$

$p \in S : x(p) = x^* \rightarrow$  divide them such that  $|S_1| = |S_2| = \frac{|S|}{2}$

3. recursively perform step 1 & 2 on  $S_1, S_2$


Conquer:

•  $d = \min_{S_1 \text{ or } S_2} \text{CP distance of a pair in}$

• to get CP of

$S = S_1 \cup S_2$

consider all points



in  $\delta$ -interval  
around  $x^*$

BUT: Could be that  
we have all points in  
there

$$\frac{h}{2} \cdot \frac{h}{2} = \frac{h^2}{4}$$

now: consider only points that are  
not too far away in terms of  
the  $y$ -coordinate

$\Rightarrow$   $\delta \times 2\delta$  rectangle


$\Rightarrow$  at most 6 points  
in there

$\Rightarrow$  constant number of comp.  
per node in  $S_n$

$\Rightarrow O(n)$

$\rightarrow$  for efficient access to the  
candidate nodes, we need them  
pre-sorted by  $y$ -coordinate  
 $\hookrightarrow$  on demand sorting takes too  
long  $O(\frac{n}{2} \log \frac{n}{2})$ , but bottom  
up merge sort can be  
applied

$\Rightarrow O(n \log n)$



Deterministic Lower Bound

↳ Sweep line and Divide & Conq.  
are asymptotically optimal  
because there is a matching  
lower bound

$\Rightarrow$  Element Uniqueness Problem (EU)

$$S = \{x_1, \dots, x_n\} \quad x_i \in \mathbb{R}$$

are all numbers unique?

$\Rightarrow$  lower bound of  $O(n \log n)$

↳ Reduction: EU can be reduced  
to CP


For each  $x_i \in S$ , we construct a point  $p = (x_i, x_i) \in \mathbb{R}^2 \Rightarrow O(n)$

( $\hookrightarrow$ ) involve CP alg. if CP has a distance of 0  $\rightarrow$  EU false  
else EU true

$\Rightarrow$  if a det. alg. solves CP faster than  $O(n \log n)$ , EU could be solved faster  $\Downarrow$

## Randomized Incremental Algorithm runs in $O(n)$

Let  $p_1, p_2, \dots, p_n$  the points in some order. Let's assume for a moment that we know about the CP distance for the points in  $p_1, \dots, p_{i-1}$ , we call this distance  $d_i$ .



build grid layer with cell length  $\delta$   
 sort all  $p_{i-1} \dots p_{i-1}$  in the respective  
 cell  $O(i-n)$

Now consider  $p_i$  and let  $c_{ae}$   
 be its grid cell, a candidate  
 to form a CP with  $p_i$  has to  
 be in  $c_{ab}$   $a \in \{e-1, e, e+1\}$   
 $b \in \{e-1, e, e+1\}$

9 cells



$\hookrightarrow$  per cell at most 4 points  
 can lie inside, bc cause  
 otherwise their distance would  
 smaller than  $\delta$   $\hookrightarrow$  at most

36 comparisons

$$\Rightarrow \text{total runtime } \sum_{i=3}^n (O(i) + O(n)) = O(n^2)$$

BUT: only have to recalc grid layer if  $p_i$  forms new CP


**EXERCISE** Fix an instance of CP and an order of the points, such that the grid needs to be rebuilt in every iteration.



Now consider the points in random order.

**EXERCISE** If points are in random order, what is the probability of point  $p_i$  to form a new CP?

$\leq \frac{2}{i}$  because 2 of  
 the  $i$  points form a CP, and  
 with every permutation being equally  
 likely the chance that  $p_i$  is one of them  
 is  $\frac{2}{i}$



Expected cost per iteration

$$O(i) \cdot \frac{2}{i} + \frac{i-2}{i} \cdot O(n)$$



$$= O(n)$$

total runtime:


$$T(n) = \sum_{i=3}^n O(n) = O(n^2)$$



EXERCISE What is the runtime of  
this CP alg. in  $d$  dimensions?



$$3^2 \cdot 2^2 = 36$$



$$3^D \cdot 2^D = 6^D$$

for fixed  $D \Rightarrow$  constant